

## Article

# Geographical–Historical Analysis of the Herbarium Specimens Representing the Economically Important Family *Amaranthaceae* (*Chenopodiaceae*-*Amaranthaceae* Clade) Collected in 1821–2022 and Preserved in the Herbarium of the Jagiellonian University in Krakow

Agata Stadnicka-Futoma <sup>1,\*</sup> and Marcin Nobis <sup>2</sup>

<sup>1</sup> Department of Soil Science, Environmental Chemistry and Hydrology, Institute of Agricultural Sciences, Environment Management and Protection, College of Natural Sciences, University of Rzeszów, ul. Zelwerowicza 8b, 35-601 Rzeszów, Poland

<sup>2</sup> Institute of Botany, Faculty of Biology, Jagiellonian University, Gronostajowa 3, 30-387 Kraków, Poland; m.nobis@uj.edu.pl

\* Correspondence: astadnicka@ur.edu.pl

**Simple Summary:** The digitization of herbarium collections is an important process that allows access to sometimes huge amounts of data that can be used in various natural sciences. In recent years, the collections of one of the thirty oldest herbariums in the world—the herbarium of the Jagiellonian University—have been digitized. This paper presents the resources of the economically very important *Amaranthaceae* family, which include 8801 herbarium sheets. They were analyzed in taxonomic, geographical, historical, and functional terms.

**Citation:** Stadnicka-Futoma, A.; Nobis, M. Geographical–Historical Analysis of the Herbarium Specimens Representing the Economically Important Family *Amaranthaceae* (*Chenopodiaceae*-*Amaranthaceae* Clade) Collected in 1821–2022 and Preserved in the Herbarium of the Jagiellonian University in Krakow. *Biology* **2024**, *13*, 435. <https://doi.org/10.3390/biology13060435>

Academic Editors: Frank H. Hellwig and Zed Rengel

Received: 1 May 2024

Revised: 31 May 2024

Accepted: 11 June 2024

Published: 13 June 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Herbaria constitute a form of documentation, store and secure comparative material, as well as constitute an extra original gene bank. They are an invaluable database among others for the biological, ethnobotanical and agricultural sciences. The digitization of herbarium collections significantly facilitates access to archival materials; however, searching them is still time-consuming. Therefore, our work aims to analyze the herbarium collection of 8801 sheets for specimens representing the economically important family *Amaranthaceae* (*Chenopodiaceae*-*Amaranthaceae* clade) deposited the oldest herbarium in Poland, the herbarium of the Jagiellonian University (KRA). These specimens have been collected from almost all the continents in dozens of countries for over 200 years. The analyses conducted, including the taxonomic coverage, geographical characteristics and origin, temporal coverage and utility importance of representative species, present the discussed resources in a more accessible way and may become a more attractive form for scientists potentially interested in more advanced research work.

**Keywords:** *Amaranthaceae*; *Chenopodiaceae*; herbarium collection; KRA; vascular plant; taxonomy; biodiversity

## 1. Introduction

Herbaria, have become increasingly popular. They constitute a form of documentation, store and secure comparative material, as well as constitute an extra original gene bank. Currently, dried plant specimens are commonly used in taxonomic, phytogeographic and cytological research. On their basis, morphometric tests can be performed, which can later be used for statistical analysis (so-called numerical taxonomy) or to describe the morphology and anatomy of individual structures at the micro level.

Taxonomists are often able to use digitized collections to identify and annotate specimens [1], to indicate typical specimens and to typify names of species for which the type has not been designated or is regarded as lost or destroyed. In recent years, publications including typification made based on the available online herbarium databases are becoming more and more common [2–10]. It is worth noting that herbaria are a source of new species, collected and often set aside as unidentified species. Following Bebbler et al. [11], about 70,000 species are still waiting to be described. In the era of digitization, the use of scans or photographs of herbarium specimens greatly facilitates the work as well as significantly saving time [9,12–14]. An important aspect is also the studies of phylogenetic relationships. Increasingly, phylogenetic analyses are based on samples collected from herbarium materials, e.g., [15–22], despite problems related to post-mortem DNA damage [23,24]. Herbaria also play an important role in research related to the distribution of plants. At the same time, they allow for analyzing changes in plant cover over many years and are used to establish phytogeographic patterns. Herbarium specimens can be used to analyze phytogeographic phenomena with temporal and spatial dynamics. On their basis, for example, the migration of invasive plants is studied, e.g., [25–31]. They have similar importance in the case of rare plants, including endangered ones, but also utility plants and weeds. They therefore constitute a biodiversity database and may be indirectly helpful in its protection. Herbarium specimens allow the identification of priority places for nature conservation, e.g., [32,33]. As it turns out, herbaria can be a source of viable diaspores. In recent years, attempts have been made to use the seeds of herbarium specimens, and the success of these studies is essential in the case of endangered or even extinct species [34]. Preserved tissues can also be used for this purpose [35,36]. Based on the data contained on the labels of herbarium specimens, it is possible to reconstruct the phenology of plant species in large areas. Phenological data are, therefore, an important tool for studying the impact of climate change on living organisms [37], particularly those related to global warming [38–42]. Based on the collected specimens or collections, it is also possible to trace the history of botanists, e.g., in relation to the expeditions they undertook, which were previously unknown [36,43]. They are also a source of information for ethnobotanists, e.g., [44].

The herbarium of the Jagiellonian University is one of the thirty oldest herbaria in the world [43]. Currently, it contains collections of vascular plants (approx. 800,000 sheets), bryophytes (25,000 specimens), fungi and slime molds (over 70,000 specimens), lichens (approx. 100,000) and algae (approx. 5000). In addition, collections of cones and fruits numbering over 1000 specimens are deposited there. In total, there are approximately one million specimens from all over the world. In 2019, the digitization of the herbarium's sheets began as part of the IMBIO project [45,46]. During this time, several large and numerous families of vascular plants were digitized, including one of the most useful and economically important families, *Amaranthaceae*.

Following the APG treatment, the family *Amaranthaceae* is represented by ca. 165 genera, comprising over 2050 species [47]. However, particular members of the family are taxonomically difficult and require integrative studies. It is worth admitting that only in the last decade have a dozen species within the family been described, e.g., [48–54].

*Amaranthaceae* contains many economically important crop plants, adapted to a wide range of climatic conditions due to the type of photosynthesis carried out, C4 types, but also due to their high nutrient content, e.g., [55–59]. For example, within the genus *Chenopodium*, there are many species, for instance, *Ch. quinoa* Willd., that contains proteins composed of amino acids that are usually missing from cereal grains (e.g., large amounts of lysine), which are considered to be richer in nutrients than cereals [60–62]. The genus *Amaranthus* is also known for containing valuable nutrients, a high protein and mineral content even compared to rice or corn grains, and an appropriate amount of exogenous amino acids (including, like the above-mentioned quinoa, lysine) [63]. Most species are also used in medicine. They have numerous medicinal properties, including anti-inflammatory, antirheumatic, diuretic, and in the case of some of them, e.g., *Achyranthes aspera* L., anticancer

activity has been demonstrated, e.g., [64–68]. Many studies indicate that, for example, *Atriplex lindleyi* Moq. may contribute to the fight against malaria and be useful in the prevention of *Candida albicans* (C.-P. Robin) Berkhout and *Pseudomonas aeruginosa* (Schröter) Migula [69]. Many species are also used in animal breeding, e.g., species of the genus *Amaranthus* are used in the production of animal feed [70]. Species used in the process of recultivation of contaminated areas are also of particular importance, e.g., *Amaranthus dubius* Mart. ex Thell. (phytoremediation of solid municipal waste landfills) [71–73], whose properties can be used to treat soil and increase the crop area. The interest in the agricultural or pharmaceutical use of this group of plants certainly results from the fact that they are also often used for food, medical, cosmetic, textile, construction, religious and even magical purposes in various parts of the world. They are therefore a rich source of research for ethnobotanists [74–78]. Because the family *Amaranthaceae* is such an economically important group of flowering plants, in this work, we decided to analyze the species deposited in the KRA herbarium in order to make the digitized collection available to a wide audience. The main aim of our work was thus the analysis of all of the herbarium sheets with representatives of the above mentioned family, in terms of the taxonomy, the origin of the specimens and the time of their acquisition, which will significantly facilitate more advanced research work. In order to highlight the importance of the *Amaranthaceae* family, the literature has been reviewed. Endangered and rare species, as well as those of economic importance, were indicated.

## 2. Materials and Methods

The subjects of this article are specimens of species from the *Amaranthaceae* family (*Chenopodiaceae-Amaranthaceae* clade) deposited in the KRA Herbarium. During digitization, the information on the herbarium labels was recorded in the appropriate columns of the Jagiellonian University's MUZUJ database. Moreover, especially in the case of the genus *Chenopodium*, the authors verified the correctness of the identification. A complete list of the specimens studied is presented in Table S1.

The families currently included in the *Amaranthaceae-Chenopodiaceae* clade have long been considered closely related, e.g., [79]. Kadereit et al. [80] listed several common features in terms of the morphology and anatomy, confirming the close relationship of species representing both families. Currently, representatives of both families are grouped into the family *Amaranthaceae*, treating *Chenopodiaceae* as its synonym [81–83]. Despite this, both Kadereit et al. [80], as well as many other researchers, e.g., [84–88], suggest that the concept of combining both families has no solid phylogenetic basis and only reflects the fact that they form a monophyletic group [80,84,86,89–91]. Therefore, they treat the families in question as the *Chenopodiaceae-Amaranthaceae* clade. Our work is based on a systematic division according to the applicable APG IV [83], while the nomenclature follows the Word Flora Online [92]. If names were not found in the mentioned list, the Plants of the World Online [93] platform was used. Information regarding the life forms of the analyzed plants or from publications cited in the text was also obtained from both sites.

## 3. Results

### 3.1. General Information


The total number of sheets with specimens representing the *Amaranthaceae* family is 8801 (Table S1). Almost 97% of them are treated in the rank of species and 6% as subspecies or variety. Slightly over 2% are identified as genera, and 0.15% are putative hybrids. The remaining 0.5% constitute specimens with illegible names, belong to the unresolved name category or are assigned to only the family. The geographical location is specified for approximately 92% of the specimens. 81% of the labels contain information on the habitat, while the remainder have no such information or the text is illegible. Generally, only a small part of the sheets provide data such as the geographical coordinates, altitude, inclination, exposure or ATPOL square number (in the case of specimens collected in Poland)

[94]. Moreover, 92% of the sheets have the collector's name and/or surname legible, while the remaining personal details are illegible or non-existent. The collection date is specified for 94% of the specimens, with full dates appearing on 86% of the specimens. In other cases, they consist of only the year, the year and the month, or are illegible. The rest are sheets without a collection date assigned.

### 3.2. Taxonomic Coverage

The specimens stored in the herbarium of the Jagiellonian University and representing the analyzed family belong to 346 species and constitute approximately 17% of the world's resources [93]. The species with the most significant number of sheets (11% of resources) is *Chenopodium album* L., while 131 species are represented by single specimens.

Generally, species representing the *Amaranthaceae-Chenopodiaceae* clade belong to 83 genera (Figure 1), constituting 47% of the world's resources [93]. The largest number of deposited specimens represent *Chenopodium* (3385 sheets), *Atriplex* (1905 sheets), and *Amaranthus* (1334 sheets). The remaining genera are represented by less than 100 sheets, and 53 genera are represented by only 1 sheet. *Atriplex*, *Salsola*, *Chenopodium* and *Amaranthus* are represented by the greatest numbers of species, while 41 genera are represented by only 1 species.




**Figure 1.** Number of species in the richest genera of the family *Amaranthaceae* in the KRA herbarium (own work).

### 3.3. Life Forms

Most specimens representing the *Amaranthaceae* family constitute herbaceous plants (77% of annuals and 23% of perennials). Shrubs constitute 22%.


### 3.4. Geographical Characteristics and Origin of Specimens

The specimens representing the *Amaranthaceae* family deposited in the KRA herbarium come from almost all over the world (Figure 2).



**Figure 2.** The origin of the specimens preserved in the KRA herbarium (own work based on the WMS EnviroSolution—<https://www.envirosolutions.pl/> (accessed on 20 January 2024)).

The country where the individual plant was collected is specified on 8069 labels. The vast majority of specimens were collected in Europe (88.11%). The percentage of plant origin in relation to individual continents is presented in Figure 3.



**Figure 3.** Percentage share of specimens representing the family *Amaranthaceae* collected on different continents and preserved in KRA (own work).

Within Europe, the vast majority of specimens (almost 6000) have been collected in Poland, 306 in Ukraine, and 127 sheets in Spain, mostly from the Canary Islands. The Asian specimens representing the family *Amaranthaceae* mainly come from Tajikistan (103 sheets), including the only specimen of *Corispermum papillosum* (Kuntze) Iljin—a species

with a relatively narrow range; 72 specimens come from Russia, with almost 67% collected between 1843 and 1933, while a single specimen of *Atriplex gmelinii* C. A. Mey. ex Bong. was collected in South Korea. The African *Amaranthaceae* flora are represented by 143 specimens. The greatest number (47) was collected from Kenya (with 46 specimens collected by A. Starzeński in the years 1943–1956). A total of 180 specimens come from North America; however, most of them (135 sheets) were collected in the United States. Single specimens have been collected in Honduras and Jamaica, including the sole specimen of *Froelichia interrupta* (L.) Moq. South America is represented by 23 specimens, collected mainly in Uruguay in the years 1899–1986. A significant part of this collection represents the genera of *Gomphrena*, *Alternanthera* and *Celosia*. A single sheet with specimens of *Iresine angustifolia* Euphrasén comes from Argentina, and four specimens of other species (including the only specimens from *Nitrophila occidentalis* (Moq.) S. Watson, *Atriplex imbricata* (Moq.) D. Dietr., *Atriplex atacamensis* Phil. come from Chile. Two sheets of *Suaeda altissima* (L.) Pall. were collected in Peru by A. Junge from the same place (one in 1907 and the other in 1909).

An important part of the collection comprises rare and endemic plant species with a limited distribution range. In the KRA herbarium, they are represented by 13 species of which the majority occur in Australia (Table S2).

### 3.5. Temporal Coverage


Specimens of vascular plants have been collected and preserved in the KRA herbarium since approximately 1750. However, the oldest collection representing the family *Amaranthaceae* (complete gathering and properly labeled), *Gomphrena globosa* L., came from 1843 and is very well-preserved (Figure 4).

Generally, the collection can be divided into three historical periods. In the first period, which took place between 1821 and 1899, 281 specimens were collected, with the largest number of specimens (38) collected in 1876. The largest number of sheets with specimens of *Amaranthaceae* was collected and deposited in KRA by A. Śleńdziński (56 sheets). They come from various regions of Ukraine. Most of the plants collected by Śleńdziński can be classified as rather common species, mostly representing the genera of *Atriplex*, *Chenopodium* and *Polycnemonum*. The most frequently collected species were *Atriplex rosea* L. (23 sheets) and *Polycnemonum arvense* L. (22 sheets). A total of 4675 specimens come from the period between 1900 and 1999. The greatest number of sheets was collected by H. Trzcińska-Tacik, 689 specimens, most of them from Poland. A small part represents Czech, Slovak and Hungarian flora. A significant number was also provided by P. Szotkowski (233) from Upper Silesia, D. Fijałkowski (220) from the Lublin region, A. Śleńdziński (185) from Ukraine, A. Sendek (142) from Silesia, J. Kornaś (120) from different countries but mostly from Poland and Zambia, K. Towpasz (106) from south-eastern Poland (mainly from the foothills) and M. Wajda (106) mainly from southern Poland. The largest number of sheets contains specimens of the following species: *Chenopodium album* L. (462), *Atriplex patula* L. (405), *Atriplex prostrata* Boucher ex DC. (274), *Chenopodium polyspermum* L. (264) and *Amaranthus retroflexus* L. (257). Moreover, 89 individual species sheets and a specimen of the genus *Cyphocarpa* come from this period. However, 3155 specimens were collected in the last and shortest period, between 2000 and 2022 (Figure 5). Most of them were collected in 2003 (699 sheets) and in 2002 (401 sheets). Over 200 sheets were collected in 2004 and each year between 2006 and 2009. The most significant numbers of specimens were collected by the following botanists: M. Nobis (373 sheets) from Poland, Tajikistan, Russia and Kyrgyzstan, as well as by M. Zarzyka-Ryszka (368 sheets), R. Piwowarczyk (252 sheets), A. Nobis (240 sheets), A. Trojcka-Brzezińska (188 sheets) and M. Bielecki (153 sheets) from Poland but from different regions. In general, 8249 labels share the collector's name for a given plant. On 144 labels, the names are partially or completely illegible, while on the remaining labels, the names of the collectors are missing. This extensive collection of *Amaranthaceae* specimens is the result of almost 1000 researchers, who within over 200 years collected the plants from different regions of the

world. The significant increase in the number of sheets collected after 1950 is related to the more intensive scientific activity of mainly employees (research trips) but also doctoral students (floristic studies) associated with the Jagiellonian University.



**Figure 4.** *Gomphrena globosa* L., one of the oldest specimens of the *Amaranthaceae* family preserved in the KRA herbarium (scan of the sheet number UJ-KRA-P-36872-N owned by the herbarium KRA).



**Figure 5.** Number of sheets with specimens from the family *Amaranthaceae* collected in individual decades (1821–2022) and preserved in KRA (own work).

### 3.6. Utility Importance of Species Representing the *Amaranthaceae* Family

Of the 346 species representing the *Amaranthaceae* family in the KRA herbarium, at least 65 have been shown to be of significant economic importance. Almost 74% have healing properties for various diseases, and some even have anti-cancer properties. A total of 25 species belonging to the genera *Atriplex* (14) and *Amaranthus* (10) are used as food. In the case of the latter, the seeds are particularly important as they are treated as “pseudo-cereals” rich in valuable nutrients [63]. These plants also provide food for animals. Some, e.g., *Aerva javanica* (Burm.f.) Schult. or *Arthrocnemum macrostachyum* (Moric.) K.Koch can be used in phytoremediation and soil reclamation [95,96]. *Alternanthera caracasana* Knuth is a potential source of bioethanol for fuel production [97]. Some species, such as the representatives of the genus *Dysphania* R. Br., are used as moth repellent and treatment for respiratory illness, and as apotropaic and insect repellents, which are blessed during Catholic church holidays [98,99]. The list of useful species representing the family *Amaranthaceae* that are preserved in the herbarium of KRA is provided in Table S3.

## 4. Discussion

Plants play a significant role in human life. They have also been an object of interest in many fields of science, and fresh or dry parts of plants are a source of valuable and reliable data. Herbaria preserving a collection of dry plants constitute a rich resource of biological diversity. In addition to their museum, historical and scientific value, they protect comparative material and constitute a specific gene bank whose importance, together with the development of molecular technologies, is increasingly appreciated [36,100]. It seems that the most important task of herbaria is to store herbarium specimens as comparative materials for taxonomic research or studies related to the analysis of plant distribution. However, the role of herbaria is much more complex, and those mentioned above are only a few examples.

Digital collections have become an urgent necessity in today’s large-scale biodiversity research, and indeed, in recent years, an intensive digitization process has been taking place in many herbaria [101]. This makes the collections available to a wide range of recipients. Easily accessible and searchable data on herbarium specimens on the Internet have greatly increased the efficiency of research in the last decade. Time previously spent travelling to collections or waiting to borrow specimens can instead be spent on data collection [9].



Analyses such as ours, as included in this article, contribute to a more accessible presentation of herbarium resources of species representing the *Amaranthaceae* family, which is of great economic importance. We believe that this way of presenting them can encourage researchers to use data from the KRA herbarium for various types of scientific research, such as ethnobotanical, pharmacological, phytogeographical or taxonomical research, for instance, related to one of the species-rich, morphologically variable and taxonomically problematic genera such as *Chenopodium*, which we intend to address in our further research. An accessible analysis, combined with the attached list of deposited perches and online access to the herbarium sheets, which are or will soon be published in the GIBF [102], provides new research opportunities and allows researchers to save time previously spent on the revision of herbarium materials directly in the place where they are stored.

## 5. Conclusions

In this work, we described the whole set of 8801 herbarium specimens representing the economically important family *Amaranthaceae*, which were collected over 200 years and are preserved in the herbarium of the Jagiellonian University (KRA). The analyses presented here, which are related to the taxonomic coverage, geographical characteristics and origin, temporal coverage, or utility importance of species, allow for a better understanding and contribute to faster familiarization with a huge body of material, the verification of which in its raw form would require a tremendous amount of work and time. In the era of increasing possibilities of using herbarium plants, the collected resources of specimens from the discussed family may become an important source for scientific studies involving many fields of natural sciences. It is worth emphasizing that individual representatives of the family *Amaranthaceae* are regarded as taxonomically problematic and can cause many problems in terms of proper identification. Therefore, the data presented in both descriptive and tabular form will be useful for comparative analyses by professional scientists conducting more advanced research works as well as by students and amateur botanists dealing with plants of the family *Amaranthaceae*.

**Supplementary Materials:** The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/biology13060435/s1>, Table S1: Database of herbarium sheets of the *Amaranthaceae* family deposited in the KRA herbarium; Table S2: A list of rare and endemic species representing the family *Amaranthaceae* and deposited in the herbarium of KRA; Table S3: List of economically useful species based on herbarium specimens of the *Amaranthaceae* family deposited in the KRA herbarium. References [59–63,65–67,69,70,72,73,93,95,96,103–163] are cited in the supplementary materials.

**Author Contributions:** Conceptualization, A.S.-F. and M.N.; methodology, A.S.-F. and M.N.; validation, A.S.-F. and M.N.; formal analysis, A.S.-F.; investigation, A.S.-F.; resources, M.N.; data curation, A.S.-F. and M.N.; writing—original draft preparation, A.S.-F.; writing—review and editing, M.N.; visualization, A.S.-F.; supervision, A.S.-F. and M.N. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding, however, M. Nobis's research was partially funded by the Institute of Botany, the Jagiellonian University, Kraków (N18/DBS/000002).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data presented in this study are available in the supplementary materials.

**Acknowledgments:** We would like to express our gratitude to the KRA herbarium for making the collection available for this study.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Harris, K.M.; Marsico, T.D. Digitizing specimens in a small herbarium: A viable workflow for collections working with limited resources. *Appl. Plant Sci.* **2017**, *5*, 1600125. <https://doi.org/10.3732/apps.1600125>.
2. Alves-Araújo, A.; dos Santos Moraes, Q.; Nichio-Amaral, R.; Miranda, V.S. Typifications in neotropical *Sapotaceae*. *PhytoKeys* **2020**, *170*, 45–69. <https://doi.org/10.3897/phytokeys.170.54718>.
3. Boltenkov, E.V.; Güner, A. Typification of some *Oncocyclus* (*Iris*, *Iridaceae*) names related to the Turkish flora. *Phytotaxa* **2020**, *468*, 045–061. <https://doi.org/10.11646/phytotaxa.468.1.2>.
4. Dalastra, C.H.; Heiden, G. Typifications of five names in *Agarista* (*Ericaceae*, *Vaccinioideae*, *Lyonieae*). Embrapa Clima Temperado—Artigo em periódico indexado (ALICE). *Phytotaxa* **2020**, *474*, 179–184. <https://doi.org/10.11646/phytotaxa.474.2.8>.
5. Kottekkattu, T.; Pradeep, A.K. Lectotypification and a new record of the genus *Eragrostiella* (*Tripogoninae*, *Cynodonteae*, *Chloridoideae*, *Poaceae*) from India. *Phytotaxa* **2020**, *464*, 109–115. <https://doi.org/10.11646/phytotaxa.464.1.10>.
6. Liu, B.B.; Liu, G.N.; Hong, D.Y.; Wen, J. Typification of 23 names in *Eriobotrya* (*Maleae*, *Rosaceae*). *PhytoKeys* **2020**, *139*, 99–118. <https://doi.org/10.3897/phytokeys.139.47967>.
7. Naive, M.A.K.; Sanders, T. Typification of two species names in *Dendrochilum* (*Orchidaceae*, subgenus *Platyclinis*). *Phytotaxa* **2020**, *470*, 298–299. <https://doi.org/10.11646/phytotaxa.470.4.4>.
8. Nobis, M.; Gudkova, P.D.; Nowak, A.; Sawicki, J.; Nobis, A. A synopsis of the genus *Stipa* (*Poaceae*) in Middle Asia, including a key to species identification, an annotated checklist, and phytogeographic analyses. *Ann. Mo. Bot. Gard.* **2020**, *105*, 1–63. <https://doi.org/10.3417/2019378>.
9. Nobis, M.; Klichowska, E.; Wolanin, M.; Nobis, A.; Nowak, A. Typification of five plant names described based on specimens collected by Józef Warszewicz in Central and South America. *PhytoKeys* **2022**, *192*, 45. <https://doi.org/10.3897/phytokeys.192.78409>.
10. Wolski, G.J.; Faltyn-Parzyńska, A.; Proćków, J. Lectotypification of the name *Stereodon nemoralis* Mitt. (*Plagiotheciaceae*), a basionym of *Plagiothecium nemorale* (Mitt.) A. Jaeger. *PhytoKeys* **2020**, *155*, 141–153. <https://doi.org/10.3897/phytokeys.155.51469>.
11. Bebb, D.P.; Carine, M.A.; Wood, J.R.I.; Wortley, A.H.; Harris, D.J.; Prance, G.T.; Davidse, G.; Paige, J.; Pennington, T.D.; Robson, N.K.B.; et al. Herbaria are a major frontier for species discovery. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 22169–22171. <https://doi.org/10.1073/pnas.101184110>.
12. Corney, D.P.A.; Clark, J.Y.; Tang, H.L.; Wilkin, P.; Corney, D.P.A.; Clark, J.Y.; Tang, H.L.; Wilkin, P. Automatic extraction of leaf characters from herbarium specimens. *Taxon* **2012**, *61*, 231–244. <https://doi.org/10.1002/tax.611016>.
13. Weaver, W.N.; Smith, S.A. From leaves to labels: Building modular machine learning networks for rapid herbarium specimen analysis with LeafMachine2. *Appl. Plant Sci.* **2023**, *11*, e11548. <https://doi.org/10.1002/aps3.11548>.
14. Carranza-Rojas, J.; Goeau, H.; Bonnet, P.; Mata-Montero, E.; Joly, A. Going deeper in the automated identification of Herbarium specimens. *BMC Evol. Biol.* **2017**, *17*, 181. <https://doi.org/10.1186/s12862-017-1014-z>.
15. Inderbitzin, P.; Lim, S.R.; Volkmann-Kohlmeyer, B.; Kohlmeyer, J.; Berbee, M.L. The phylogenetic position of *Spathulospora* based on DNA sequences from dried herbarium material. *Mycol. Res.* **2004**, *108*, 737–748. <https://doi.org/10.1017/s0953756204000206>.
16. Zuntini, A.R.; Fonseca, L.H.M.; Lohmann, L.G. Primers for phylogeny reconstruction in *Bignoniaceae* (*Bignoniaceae*) using herbarium samples. *Appl. Plant Sci.* **2013**, *1*, 2–5. <https://doi.org/10.3732/apps.130001>.
17. Besnard, G.; Christin, P.A.; Malé, P.J.G.; Lhuillier, E.; Lauzeral, C.; Coissac, E.; Vorontsova, M.S. From museums to genomics: Old herbarium specimens shed light on a C3 to C4 transition. *J. Exp. Bot.* **2014**, *65*, 6711–6721. <https://doi.org/10.1093/jxb/eru395>.
18. Muñoz-Rodríguez, P.; Carruthers, T.; Wood, J.R.I.; Williams, B.R.M.; Weitemier, K.; Kronmiller, B.; Goodwin, Z.; Sumadijaya, A.; Anglin, N.L.; Filer, D.; et al. A taxonomic monograph of *Ipomoea* integrated across phylogenetic scales. *Nat. Plants* **2019**, *5*, 1136–1144. <https://doi.org/10.1038/s41477-019-0535-4>.
19. Vintsek, L.; Klichowska, E.; Nowak, A.; Nobis, M. Genetic differentiation, demographic history and distribution models of high alpine endemic vicariants outline the response of species to predicted climate changes in a Central Asian biodiversity hotspot. *Ecol. Indic.* **2022**, *144*, 109419. <https://doi.org/10.1016/j.ecolind.2022.109419>.
20. Nobis, M.; Wróbel, S.; Klichowska, E.; Nowak, A.; Wróbel, A.; Nobis, A.; Paszko, B.; Świarszcz, S.; Chen, W.; Kauzal, P.; et al. New national and regional plant records: Contribution to the flora of the Old World countries. *Acta Soc. Bot. Pol.* **2023**, *92*, 1–21. <https://doi.org/10.5586/asbp/162050>.
21. Wróbel, A.; Klichowska, E.; Nobis, M. Hybrids as mirrors of the past: Genomic footprints reveal spatio-temporal dynamics and extinction risk of alpine extremophytes in the mountains of Central Asia. *Front. Plant Sci.* **2024**, *15*, 1–22. <https://doi.org/10.3389/fpls.2024.1369732>.
22. Sinaga, P.; Klichowska, E.; Nowak, A.; Nobis, M. Hybridization and introgression events in co-occurring populations of closely related grasses (*Poaceae*: *Stipa*) in high mountain steppes of Central Asia. *PLoS ONE* **2024**, *19*, e0298760. <https://doi.org/10.1371/journal.pone.0298760>.
23. Staats, M.; Cuenca, A.; Richardson, J.E.; Vrieling-van Ginkel, R.; Petersen, G.; Seberg, O.; Bakker, F.T. DNA damage in plant Herbarium tissue. *PLoS ONE* **2011**, *6*, e28448. <https://doi.org/10.1371/journal.pone.0028448>.
24. McAssey, E.V.; Downs, C.; Yorkston, M.; Morden, C.; Heyduk, K. A comparison of freezer-stored DNA and herbarium tissue samples for chloroplast assembly and genome skimming. *Appl. Plant Sci.* **2023**, *11*, 1–10. <https://doi.org/10.1002/aps3.11527>.
25. Pyšek, P. *Heracleum mantegazzianum* in the Czech Republic: Dynamics of spreading from the historical perspective. *Folia Geobot. Phytotax.* **1991**, *26*, 439–454.

26. Pyšek, P.; Prach, K. Plant invasion and the role of riparian habitats: A comparison of four species alien to central Europe. *J. Biogeogr.* **1993**, *20*, 413–420. <https://doi.org/10.2307/2845589>.
27. Pyšek, P.; Prach, K. Invasion dynamics of *Impatiens glandulifera*: A century of spreading reconstructed. *Biol. Conserv.* **1995**, *74*, 41–48. <https://doi.org/10.1016/0006-320700013-T>.
28. Lambrinos, J.G. The expansion history of a sexual and asexual species of *Cortaderia* in California, USA. *J. Ecol.* **2001**, *89*, 88–98. <https://doi.org/10.1046/j.1365-2745.2001.00524.x>.
29. Mihulka, S.; Pyšek, P. Invasion history of *Oenothera congeners* in Europe: A comparative study of spreading rates in the last 200 years. *J. Biogeogr.* **2001**, *28*, 597–609. <https://doi.org/10.1046/j.1365-2699.2001.00574.x>.
30. Delisle, F.; Lavoie, C.; Jean, M.; Lachance, D. Reconstructing the spread of invasive plants: Taking into account biases associated with herbarium specimens. *J. Biogeogr.* **2003**, *30*, 1033–1042. <https://doi.org/10.1046/j.1365-2699.2003.00897.x>.
31. Ni, M. Herbarium records reveal multiple phases in the relationship between minimum residence time and invasion ranges of alien plant species. *Plants People Planet* **2022**, *5*, 47–57. <https://doi.org/10.1002/ppp3.10327>.
32. Hernández, H.M.; Navarro, M. A new method to estimate areas of occupancy using herbarium data. *Biodivers. Conserv.* **2007**, *16*, 2457–2470. <https://doi.org/10.1007/s10531-006-9134-6>.
33. Verspagen, N.; Erkens, R.H.J. A method for making Red List assessments with herbarium data and distribution models for species-rich plant taxa: Lessons from the Neotropical genus *Guatteria* (Annonaceae). *Plants People Planet* **2023**, *5*, 536–546. <https://doi.org/10.1002/ppp3.10309>.
34. Godefroid, S.; van de Vyver, A.; Stoffelen, P.; Robbrecht, E.; Vanderborght, T. Testing the viability of seeds from old herbarium specimens for conservation purposes. *Taxon* **2011**, *60*, 565–569. <https://doi.org/10.1002/tax.602022>.
35. Rocchetti, A.G.; Davis, C.; Caneva, G.; Bacchetta, G.; Fabrini, G.; Fenu, G.; Foggi, B.; Galasso, G.; Gargano, D.; Giusso del Galdo, G.; et al. A pragmatic and prudent consensus on the resurrection of extinct plant species using herbarium specimens. *Taxon* **2022**, *71*, 168–177. <https://doi.org/10.1002/tax.12601>.
36. Heberling, J.M.; Prather, L.A.; Tonsor, S.J. The changing uses of Herbarium data in an era of global change: An overview using automated content analysis. *BioScience* **2019**, *69*, 812–822. <https://doi.org/10.1093/biosci/biz094>.
37. Lavoie, C.; Lachance, D. A new herbarium-based method for reconstructing the phenology of plant species across large areas. *Am. J. Bot.* **2006**, *93*, 512–516. <https://doi.org/10.3732/ajb.93.4.512>.
38. Primack, D.; Imbres, C.; Primack, R.B.; Miller-Rushing, A.J.; Del Tredici, P. Herbarium specimens demonstrate earlier flowering times in response to warming in Boston. *Am. J. Bot.* **2004**, *91*, 1260–1264. <https://doi.org/10.3732/ajb.91.8.1260>.
39. Miller-Rushing, A.J.; Primack, R.B.; Primack, D.; Mukunda, S. Photographs and herbarium specimens as tools to document phenological changes in response to global warming. *Am. J. Bot.* **2006**, *93*, 1667–1674. <https://doi.org/10.3732/ajb.93.11.1667>.
40. Hart, R.; Salick, J.; Ranjitkar, S.; Xu, J. Herbarium specimens show contrasting phenological responses to Himalayan climate. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 10615–10619. <https://doi.org/10.1073/pnas.1403376111>.
41. Rawal, D.S.; Kasel, S.; Keatley, M.R.; Nitschke, C.R. Herbarium records identify sensitivity of flowering phenology of eucalypts to climate: Implications for species response to climate change. *Austral Ecol.* **2014**, *40*, 117–125. <https://doi.org/10.1111/aec.12183>.
42. Davis, C.C.; Willis, C.G.; Connolly, B.; Kelly, C.; Ellison, A.M. Herbarium records are reliable sources of phenological change driven by climate and provide novel insights into species' phenological cueing mechanisms. *Amer. J. Bot.* **2015**, *102*, 1599–1609. <https://doi.org/10.3732/ajb.1500237>.
43. Plewa, A.; Köhler, P. *Pinaceae* in the herbarium of the Institute of botany at the Jagiellonian University, Kraków, Poland (KRA). *Acta Soc. Bot. Pol.* **2019**, *88*, 1–13. <https://doi.org/10.5586/asbp.3624>.
44. Azab, A. *Amaranthaceae* plants of Israel and palestine: Medicinal activities and unique compounds. *Eur. Chem. Bull.* **2020**, *9*, 366–400. <https://doi.org/10.17628/ecb.2020.9.366-400>.
45. Tykarski, P. Natural data resources of Polish scientific institutions—variety, history, importance—introduction. *Kosmos* **2021**, *70*, 139–145. [https://doi.org/10.36921/kos.2021\\_2798](https://doi.org/10.36921/kos.2021_2798).
46. Knutelski, S.; Nobis, M.; Pyrcz, T.; Fiałkowski, W. Zasoby informacji o różnorodności biotycznej w kolekcjach przyrodniczych Uniwersytetu Jagiellońskiego. *Kosmos* **2021**, *70*, 273–289. [https://doi.org/10.36921/kos.2021\\_2786](https://doi.org/10.36921/kos.2021_2786).
47. Christenhusz, J.M.; Byng, J.W. The number of know plants species in the world and its annual increase. *Phytotaxa* **2016**, *261*, 201–217. <https://doi.org/10.11646/phytotaxa.261.3.1>.
48. Flores-Olvera, H.; Zumaya, S.; Borsch, T. Two new species of *Iresine* (Amaranthaceae: Gomphrenoideae) from Mexico supported by morphological and molecular characters. *Willdenowia* **2016**, *46*, 165–174. <https://doi.org/10.3372/wi.46.46113>.
49. Alonso, Á.M.; Crespo, M.B.; Freitag, H. *Salicornia cuscoensis* (Amaranthaceae/Chenopodiaceae), a new species from Peru (South America). *Phytotaxa* **2017**, *319*, 254–262. <https://doi.org/10.11646/phytotaxa.319.3.4>.
50. Arya, S.; Kumar, V.S.A.; Vishnu, W.K.; Kumar, T.R. *Amaranthus saradhiana* (Amaranthaceae)—a new species from southern Western Ghats of Kerala, India. *Phytotaxa* **2019**, *403*, 230–238. <https://doi.org/10.11646/phytotaxa.403.3.7>.
51. Arya, S. *Alternanthera ebracteolata* (Amaranthaceae), a new species from Kerala (SW-India). *Phytotaxa* **2021**, *480*, 191–196. <https://doi.org/10.11646/phytotaxa.480.3.7>.
52. Arya, S.; Iamonico, D.; Sánchez-Del Pino, I.; Kumar, V.N.S.A. *Alternanthera indica* (Amaranthaceae), a new species from Kerala (India). *Phytotaxa* **2021**, *482*, 191–196. <https://doi.org/10.11646/phytotaxa.482.2.7>.
53. Mosyakin, S.L.; Mandák, B. *Chenopodium ucrainicum* (Chenopodiaceae/Amaranthaceae sensu APG), a new diploid species: A morphological description and pictorial guide. *Ukr. Bot. J.* **2020**, *77*, 237–248. <https://doi.org/10.15407/ukrbotj77.04.237>.

54. Chatelain, C.; Uotila, P.; Benhouhou, S.; Mombrial, F.; Mesbah, M.; Baa, S.; Benghanem, A.N. *Chenopodium hoggarensis* (Amaranthaceae), a new species from Algeria and Chad. *Willdenowia* **2022**, *52*, 75–81. <https://doi.org/10.3372/wi.52.52104>.
55. Simmonds, N.W. The grain chenopods of the tropical American highlands. *Econ. Bot.* **1965**, *19*, 223–235.
56. Galwey, N.W.; Leakey, C.L.A.; Price, K.R.; Fenwick, G.R. Chemical composition and nutritional characteristics of quinoa (*Chenopodium quinoa* Willd.). *Human Nutr. Food Sci. Nutr.* **1990**, *42F*, 245–261. <https://doi.org/10.1080/09543465.1989.11904148>.
57. Dincă, L.; Dincă, M.; Pantea, S.D.; Timiș-Gansac, V.; Oneț, C. *Amaranthus* Plant—between myth and usage. *Natural Resources Sustain. Dev.* **2018**, *8*, 9–16. <https://doi.org/10.31924/nrsd.v8i1.002>.
58. Bajwa, A.A.; Zulfiqar, U.; Sadia, S.; Bhowmik, P.; Chauhan, B.S. A global perspective on the biology, impact and management of *Chenopodium album* and *Chenopodium murale*: Two troublesome agricultural and environmental weeds. *Environ. Sci. Pollut. Res.* **2019**, *26*, 5357–5371. <https://doi.org/10.1007/s11356-018-04104-y>.
59. Karous, O.; Ben Haj Jilani, I.; Ghrabi-Gammar, Z. Ethnobotanical study on plant used by Semi-Nomad Descendants' Community in Ouled Dabbeb—Southern Tunisia. *Plants* **2021**, *10*, 642. <https://doi.org/10.3390/plants10040642>.
60. Schlick, G.; Bubenheim, D.L. *Quinoa*: Candidate crop for NASA's controlled ecological life support systems. In *Progress in New Crops*; Janick, J., Ed.; ASHS Press: Arlington, VA, USA, 1996; pp. 632–640.
61. Dini, I.; Schettino, O.; Simioli, T.; Dini, A. Studies on the constituents of *Chenopodium quinoa* seeds: Isolation and characterization of new triterpene saponins. *J. Agric. Food Chem.* **2001**, *49*, 741–746. <https://doi.org/10.1021/jf000971y>.
62. Iqbal, M. An Assessment of quinoa (*Chenopodium quinoa* Willd.) potential as a grain crop on marginal lands in Pakistan. *Am. Eurasian J. Agric. Environ. Sci.* **2015**, *15*, 16–23. <https://doi.org/10.5829/idosi.ajeaes.2015.15.1.12474>.
63. Aderibigbe, O.R.; Ezekiel, O.O.; Owolade, S.O.; Korese, J.K.; Sturm, B.; Hensel, O. Exploring the potentials of underutilized grain amaranth (*Amaranthus* spp.) along the value chain for food and nutrition security: A review. *Crit. Rev. Food Sci. Nutr.* **2020**, *62*, 656–669. <https://doi.org/10.1080/10408398.2020.1825323>.
64. Kokanova-Nedialkova, Z.; Nedialkov, P.T.; Nikolov, S.D. The genus *Chenopodium*: Phytochemistry, ethnopharmacology and pharmacology. *Pharmacogn. Rev.* **2009**, *3*, 280–306.
65. Donia El Raheim Mohammed, A.; Ibrahim Alqasoumi, S.; Mahmoud Radwan, A.; Burand, J.; Craker, L.E. Phytochemical screening and insecticidal activity of three plants from *Chenopodiaceae* family. *J. Med. Plant Res.* **2012**, *6*, 5863–5867. <https://doi.org/10.5897/JMPR11.1629>.
66. Soumia, H.; Zahia, B.; Aminata, O.E.H.K.; Hanane, B.; Zohra, B.; Yasmina, B.; Insaf, A.K.; Sara, M.; Fatma, S.; Meriem, T. Ethnobotanical study and phytochemical screening of six medicinal plants used in traditional medicine in the Northeastern Sahara of Algeria (area of Ouargla). *J. Med. Plant Res.* **2015**, *9*, 1049–1059. <https://doi.org/10.5897/jmpr2015.5932>.
67. Alharbi, N. Survey of plant species of medical importance to treat digestive tract diseases in Tabuk Region, Saudi Arabia. *J. King Abdulaziz Univ. Sci.* **2017**, *29*, 51–61. <https://doi.org/10.4197/sci.29-1.6>.
68. Dobrowolska, E.; Motyka, S.; Szopa, A.; Ekiert, H. *Achyranthes bidenata* (grzebyk dwuzębny)—charakterystyka botaniczna, ekologiczna, fitochemiczna oraz zastosowanie w lecznictwie. *Farmacja Polska* **2021**, *77*, 717–732. <https://doi.org/10.32383/farmpol/146835>.
69. El Souda, S.S.; Matloub, A.A.; Nepveu, F.; Valentin, A.; Roques, C. Phenolic composition and prospective anti-infectious properties of *Atriplex lindleyi*. *Asian Pac. J. Trop. Dis.* **2015**, *5*, 786–791. <https://doi.org/10.1016/S2222-180860931-8>.
70. Ganjare, A.; Raut, N. Nutritional and medicinal potential of *Amaranthus spinosus*. *RJPP* **2019**, *8*, 3149–3156.
71. Eissa, M.A. Phytoextraction mechanism of Cd by *Atriplex lentiformis* using some mobilizing agents. *Ecol. Eng.* **2017**, *108*, 220–226. <https://doi.org/10.1016/j.ecoleng.2017.08.025>.
72. Vromman, D.; Flores-Bavestrello, A.; Šlejkovec, Z.; Lapaille, S.; Teixeira-Cardoso, C.; Briceño, M.; Lutts, S. Arsenic accumulation and distribution in relation to young seedling growth in *Atriplex atacamensis* Phil. *Sci. Total Environ.* **2011**, *412–413*, 286–295. <https://doi.org/10.1016/j.scitotenv.2011.09.085>.
73. Ma, D.; He, Z.; Bai, X.; Wang, W.; Zhao, P.; Lin, P.; Zhou, H. *Atriplex canescens*, a valuable plant in soil rehabilitation and forage production. A review. *Sci. Total Environ.* **2022**, *15*, 150287. <https://doi.org/10.1016/j.scitotenv.2021.150287>.
74. Qureshi, R.; Bhatti, G.R. Folklore uses of *Amaranthaceae* family from Nara Desert, Pakistan. *Pak. J. Bot.* **2009**, *41*, 1565–1572.
75. Bieski, I.G.C.; Leonti, M.; Arnason, J.T.; Ferrier, J.; Rapinski, M.; Violante, I.M.P.; Balogun, S.O.; Pereira, J.F.; de Figueiredo, R.C.; Lopes, C.R.; et al. Ethnobotanical study of medicinal plants by population of Valley of Jurueña Region, Legal Amazon, Mato Grosso, Brazil. *J. Ethnopharmacol.* **2015**, *173*, 383–423. <https://doi.org/10.1016/j.jep.2015.07.025>.
76. Saranraj, P.; Lakshmi, B.; Suganthi, K. Ethnobotanical survey of medicinal plants from Vellore district, Tamil nadu, India. *Int. J. Adv. Res. Biol. Sci.* **2016**, *3*, 238–246. <https://doi.org/10.22192/ijarbs>.
77. Singh, G.; Kumar, J. Studies on underutilized weeds of family *Amaranthaceae* used as edibles by the Munda tribe of Jharkhand, India. *Ann. Plant Sci.* **2019**, *8*, 3495–3498. <https://doi.org/10.21746/aps.2018.8.2.1>.
78. Manzanero-Medina, G.I.; Vásquez-Dávila, M.A.; Lustre-Sánchez, H.; Pérez-Herrera, A. Ethnobotany of food plants (quelites) sold in two traditional markets of Oaxaca, Mexico. *S. Afr. J. Bot.* **2020**, *130*, 215–223. <https://doi.org/10.1016/j.sajb.2020.01.002>.
79. Kühn, U.; Bittrich, V.; Carolin, R.; Freitag, H.; Hedge, I.C.; Uotila, P.; Wilson, P.G. *Chenopodiaceae*. In *The Families and Genera of Vascular Plants*; Kubitzki, K., Rohwer, J.G., Bittrich, V., Eds.; Springer, Berlin, Germany, 1993; Volume 2, pp. 253–281.
80. Kadereit, G.; Borsch, T.; Weising, K.; Freitag, H. Phylogeny of *Amaranthaceae* and *Chenopodiaceae* and the evolution of C4 photosynthesis. *J. Plant Sci.* **2003**, *164*, 959–986. <https://doi.org/10.1086/378649>.
81. Angiosperm Phylogeny Group. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Bot. J. Linn. Soc.* **2003**, *141*, 399–436. <https://doi.org/10.1046/j.1095-8339.2003.t01-1-00158.x>.

82. Angiosperm Phylogeny Group. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.* **2009**, *161*, 105–121. <https://doi.org/10.1111/j.1095-8339.2009.00996.x>.
83. Angiosperm Phylogeny Group. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot. J. Linn. Soc.* **2016**, *181*, 1–20. <https://doi.org/10.1111/boj.12385>.
84. Brockington, S.F.; Alexandre, R.; Ramdial, J.; Moore, M.J.; Crawley, S.; Dhingra, A.; Hilu, K.; Soltis, D.E.; Soltis, P.S. Phylogeny of the *Caryophyllales* sensu lato: Revisiting hypotheses on pollination biology and perianth differentiation in the core *Caryophyllales*. *Int. J. Plant Sci.* **2009**, *170*, 627–643. <https://doi.org/10.1086/597785>.
85. Sukhorukov, A.P. *The Carpology of the Chenopodiaceae with Reference to the Phylogeny, Systematics and Diagnostics of Its Representatives*; Grif & Co.: Tula, Russia, 2014.
86. Hernández-Ledesma, P.; Berendsohn, W.G.; Borsch, T.; von Mering, S.; Akhani, H.; Arias, S.; Castañeda-Noa, I.; Eggli, U.; Eriksson, R.; Flores-Olvera, H.; et al. A taxonomic backbone for the global synthesis of species diversity in the angiosperm order *Caryophyllales*. *Willdenowia* **2015**, *45*, 281–383. <https://doi.org/10.3372/WI.45.45301>.
87. Sukhorukov, A.P.; Nilova, M.V.; Krinitsina, A.A.; Zaika, M.A.; Erst, A.S.; Shepherd, K.A. Molecular phylogenetic data and seed coat anatomy resolve the generic position of some critical *Chenopodioideae* (*Chenopodiaceae*-*Amaranthaceae*) with reduced perianth segments. *PhytoKeys* **2018**, *109*, 103–128. <https://doi.org/10.3897/phytokeys.109.28956>.
88. Sukhorukov, A.P.; Liu, P.L.; Kushunina, M. Taxonomic revision of *Chenopodiaceae* in Himalaya and Tibet. *PhytoKeys* **2019**, *116*, 1–141. <https://doi.org/10.3897/phytokeys.116.27301>.
89. Cuénoud, P.; Savolainen, V.; Chatrou, L.W.; Powell, M.; Grayer, R.J.; Chase, M.W. Molecular phylogenetics of *Caryophyllales* based on nuclear 18S rDNA and plastid *rbcL*, *atpB*, and *matK* DNA sequences. *Amer. J. Bot.* **2002**, *89*, 132–144. <https://doi.org/10.3732/ajb.89.1.132>.
90. Müller, K.; Borsch, T. Phylogenetics of *Amaranthaceae* based on *matK*/*trnK* sequence data—Evidence from parsimony, likelihood, and bayesian analyses. *Ann. Missouri Bot. Gard.* **2005**, *92*, 66–102. <https://doi.org/10.3906/bot-0707-15>.
91. Schäferhoff, B.; Müller, K.F.; Borsch, T. *Caryophyllales* phylogenetics: Disentangling *Phytolaccaceae* and *Molluginaceae* and description of *Microteaceae* as a new isolated family. *Willdenowia* **2009**, *39*, 209–228. <https://doi.org/10.3372/wi.39.39201>.
92. WFO. The World Flora Online. Available online: <https://www.worldfloraonline.org/> (accessed on 10 October 2022).
93. POWO. Plants of the World Online, Facilitated by the Royal Botanic Gardens, Kew. Available online: <https://powo.science.kew.org/> (accessed on 10 October 2022).
94. Zając, A. Założenia metodyczne “Atlasu rozmieszczenia roślin naczyniowych w Polsce”. *Wiadomości Botaniczne* **1978**, *22*, 145–155.
95. Novelty, P.; Watson, I. Successful grassland regeneration in a severely degraded catchment: A whole of government approach in north west Australia. In *Climate and Land Degradation. Environmental Science and Engineering Series*; Chapter 26; Sivakumar, M.V.K., Ndiang’ui, N., Eds.; Springer: Berlin, Germany, 2007; pp. 469–484.
96. Megharbi, A.; Kechairi, R. Ethnobotanical characterization of halophytes with medicinal virtues, Case of the Macta wetland flora: North-West Algeria. *Genet. Biodivers. J.* **2021**, *5*, 135–145. <https://doi.org/10.46325/gabj.v5i2.82>.
97. Malthesh, S.R.Y.; Achar, R.R.; Cathrine, A.A.; Vadiraj, K.T. Extraction of alcohols from non-edible agricultural weed, lignocellulouic feedstock—*Alternanthera caracasana*. *World J. Environ. Biosci.* **2023**, *12*, 27–32. <https://doi.org/10.51847/OEAOcF6m7y>.
98. Oklejewicz, K.; Łuczaj, Ł. Plants blessed in churches on assumption day in the southern suburbs of Rzeszow with special reference to *Dysphania schraderiana* (Schult.) Mosyakin & Clemants (in polish). *Etnobiologia Polska* **2015**, *5*, 15–26.
99. Łuczaj, Ł.; Wolanin, M.; Drobnik, J.; Kujawska, M.; Dumanowski, J.; Walker, K.; Tomczyk, M. *Dysphania schraderiana* (Schult.) Mosyakin & Clemants—An overlooked medicinal and ritual plant used in Poland. *J. Ethnopharmacol.* **2022**, *284*, 114755. <https://doi.org/10.1016/j.jep.2021.114755>.
100. Lavoie, C. Biological collections in an ever changing world: Herbaria as tools for biogeographical and environmental studies. *Perspect. Pl. Ecol. Evol. Syst.* **2013**, *15*, 68–76. <https://doi.org/10.1016/j.ppees.2012.10.002>.
101. Kovtonyuk, N.; Han, I.; Gatilova, E. *Digital Herbarium Collections of the Central Siberian Botanical Garden SB RAS, Novosibirsk, Russia*; Springer Proceedings in Earth and Environmental Sciences, Springer: Cham, Switzerland, 2019; pp. 22–27. [https://doi.org/10.1007/978-3-030-11720-7\\_4](https://doi.org/10.1007/978-3-030-11720-7_4).
102. GIBF. Global Biodiversity Information Facility. Available online: <https://www.gbif.org/> (accessed on 10 October 2022).
103. Thiv, M.; Thulin, M.; Kilian, N.; Linder, H.P. Eritreo–Arabian Affinities of the Socotran flora as revealed from the molecular phylogeny of *Aerva* (*Amaranthaceae*). *Syst. Bot.* **2006**, *31*, 560–570. Available online: <https://www.jstor.org/stable/25064185> (accessed on 10 October 2022).
104. Miller, A. *Aerva microphylla*. The IUCN Red List of Threatened Species 2004: e.T44753A10946247. Available online: <https://www.iucnredlist.org/species/44753/10946247> (accessed on 10 October 2023).
105. Weakley, A.; Bucher, M.; Murdock, N. *Recovery Plan for Seabeach Amaranth (Amaranthus pumilus)*; U.S. Fish and Wildlife Service, Southeast Region: Atlanta, GA, USA, 1996; pp. 1–59.
106. Marcone, M.F. First report of the characterization of the threatened plant species *Amaranthus pumilus* (seabeach amaranth). *J. Agric. Food Chem.* **2000**, *48*, 378–382. <https://doi.org/10.1021/jf991133a>.
107. NSE. Nature Serve Explorer. Available online: [https://explorer.natureserve.org/Taxon/ELEMENT\\_GLOBAL.2.141860/Amaranthus\\_pumilus](https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.141860/Amaranthus_pumilus) (accessed on 10 Mach 2024).

108. IUNC. The IUCN Red List of Threatened Species 2004. Available online: <https://www.iucnredlist.org/species/199895/2617800> (accessed on 7 March 2024).
109. Hadjiev, V. *Anabasis eugeniae*. The IUCN Red List of Threatened Species 2014. Available online: <https://www.iucnredlist.org/species/199895/2617800> (accessed on 7 March 2024).
110. Ntore, S.; Beentje, H.J.; Fischer, E.; Kabuye, C.; Kalema, J.; Kayombo, C.; Luke, W.R.Q.; Nshutiyayesu, S. *Celosia elegantissima*. The IUCN Red List of Threatened Species 2019. Available online: <https://www.iucnredlist.org/species/103581524/103648247> (accessed on 7 March 2024).
111. Foziljonov, S. Theoretical analysis of the degree of scarcity of representatives of the family of *Chenopodiaceae* in the Feragana Valley. *Cent. Asian J. Med. Sci.* **2021**, *2*, 295–299.
112. Martínez Salas, E.; Fuentes, A.C.D.; Samain, M.-S. *Celosia monosperma*. The IUCN Red List of Threatened Species 2020. Available online: <https://www.iucnredlist.org/species/136621713/137375989> (accessed on 7 March 2024).
113. Raimondo, D.; von Staden, L.; Foden, W.; Victor, J.E.; Helme, N.A.; Turner, R.C.; Kamundi, D.A.; Manyama, P.A. *Red List of South African Plants*; *Strelitzia* 25; South African National Biodiversity Institute: Pretoria, South Africa, 2009.
114. Buira, A.; Fraga i Arquimbau, P. *Salsola papillosa*. The IUCN Red List of Threatened Species 2017. Available online: <https://www.iucnredlist.org/species/103535407/103535411> (accessed on 10 October 2022).
115. Shafique, S.; Javaid, A.; Bajwa, R.; Shafiq, S. Biological control of *Achyranthes aspera* and *Xanthium strumarium* in Pakistan. *Pak. J. Bot.* **2007**, *39*, 2607–2610.
116. Sumeet, D.; Raghvendra, D.; Kushagra, M. *Achyranthes aspera* Linn. (Chirchira): A magic herb in folk medicine. *Ethnobot. Leaflet.* **2008**, *12*, 670–676. <https://doi.org/10.7439/ijbar.v2i6.36>.
117. Varuna, K.M.; Khan, M.U.; Sharma, P.K. Review on *Achyranthes aspera*. *J. Pharm. Pract. Res.* **2010**, *3*, 714–717.
118. Ozturk, M.; Mermut, A.R.; Celik, A. Urbanisation, land use, land degradation and environment. *Ambio* **2011**, *51*, 1446–1458. <https://doi.org/10.1007/s13280-022-01701-7>.
119. Mukherjee, H.; Ojha, D.; Baga, P.; Chandelb, H.S.; Bhattacharyyad, S.; Chatterjeed, T.K.; Mukherjeed, P.K.; Chakrabortia, S.; Chattopadhyaya, D. Anti-herpes virus activities of *Achyranthes aspera*: An Indian ethnobotany, and its triterpene acid. *Microbiol. Res.* **2013**, *168*, 238–244. <https://doi.org/10.1016/j.micres.2012.11.002>.
120. Hasan, S. Pharmacological and medicinal uses of *Achyranthes aspera*. *Int. J. Sci. Environ.* **2014**, *3*, 123–129.
121. Singh, N.; Mrinal, P.S.; Gupta, V.K. A review on pharmacological aspects of *Achyranthes aspera*. *Int. J. Pharmacogn. Chin. Med.* **2019**, *3*, 000188. <https://doi.org/10.23880/ipcm-16000188>.
122. Movaliya, V.; Zaveri, M. A review on the pashanbheda plant “*Aerva javanica*”. *Int. J. Pharm. Sci. Rev. Res.* **2014**, *25*, 268–275.
123. Suleiman, M.H.A. Ethnobotanical, phytochemical and biological study of *Tamarix aphylla* and *Aerva javanica* medicinal plants growing in the Asir Region, Saudi Arabia. *Trop. Conserv. Sci.* **2019**, *12*, 1940082919869480. <https://doi.org/10.1177/1940082919869480>.
124. Soliman, M.A. Cytogenetical studies on *Aerva javanica* (*Amaranthaceae*). *Fl. Medit.* **2006**, *16*, 333–339.
125. Rahman, A.H.M.M.; Gulshana, M.I.A. Taxonomy and medicinal uses on *Amaranthaceae* family of Rajshahi, Bangladesh. *Appl. Ecol. Environ. Res.* **2014**, *2*, 54–59. <https://doi.org/10.12691/aees-2-2-3>.
126. Qureshi, R.; Raza Bhatti, G. Ethnobotany of plants used by the Thari people of Nara Desert, Pakistan. *Fitoterapia* **2008**, *79*, 468–473. <https://doi.org/10.1016/j.fitote.2008.03.010>.
127. Zhao, P.; Li, X.; Sun, H.; Zhao, X.; Wang, X.; Ran, R.; Zhao, J.; Wei, Y.; Liu, X.; Chen, G. Healthy values and de novo domestication of sand rice (*Agriophyllum squarrosum*), a comparative view against *Chenopodium quinoa*. *Crit. Rev. Food. Sci. Nutr.* **2023**, *63*, 4188–4209. <https://doi.org/10.1080/10408398.2021.1999202>.
128. Qian, C.; Yin, H.; Shi, Y.; Zhao, J.; Yin, C.; Luo, W.; Dong, Z.; Xia Yan, G.C.; Wang, X.-R.; Ma, X.-F. Population dynamics of *Agriophyllum squarrosum*, a pioneer annual plant endemic to mobile sand dunes, in response to global climate change. *Sci. Rep.* **2016**, *6*, 26613. <https://doi.org/10.1038/srep26613>.
129. Genievskaya, Y.; Abugaliev, S.; Zhubanysheva, A.; Turuspekov, Y. Morphological description and DNA barcoding study of sand rice (*Agriophyllum squarrosum*, *Chenopodiaceae*) collected in Kazakhstan. *BMC Plant Biol.* **2017**, *17*, 177. <https://doi.org/10.1186/s12870-017-1132-1>.
130. Yin, X.; Yan, X.; Qian, C.; Zhou, S.; Fang, T.; Fan, X.; Gao, Y.; Chang, Y.; Yang, J.; Ma, X.F. Comparative transcriptome analysis to identify genes involved in terpenoid biosynthesis in *Agriophyllum squarrosum*, a folk medicinal herb native to Asian temperature deserts. *Plant Biotechnol. Rep.* **2021**, *15*, 369–387. <https://doi.org/10.1007/s11816-021-00674-5>.
131. Dhanya, V.; Ragavendran, U.; Aathira, M.; Saipriya, V.; Priya, M.S.; Palanisamy, S.; Siddhraj, P. Biochemical composition and antioxidant potential of raw and hydrothermal treated of two underutilized leafy vegetables *Amaranthus dubius* and *Allmania nodiflora*. *Int. J. Food Sci. Nut.* **2017**, *2*, 95–107.
132. Banerjee, S.; Joglekar, A.; Mishra, M. A critical review on importance of green leafy vegetables. *Int. J. Appl. Sci.* **2015**, *2*, 124–132.
133. Canales-Martínez, M.; Hernández-Delgado, T.; Flores-Ortiz, C.; Durán-Díaz, A.; García-Bores, A.M.; Avila-Acevedo, G. Antimicrobial activity of *Alternanthera caracasana*. *Pharm. Biol.* **2005**, *43*, 305–307. <https://doi.org/10.1080/13880200590951685>.
134. Hwong, C.S.; Leong, K.H.; Abdul Aziz, A.; Mat Junit, S.; Mohd Noor, S.; Kong, K.W. *Alternanthera sessilis*: Uncovering the nutritional and medicinal values of an edible weed. *J. Ethnopharmacol.* **2022**, *298*, 115608. <https://doi.org/10.1016/J.JEP.2022.115608>.

135. George, S.; Bhalerao, S.V.; Lidstone, E.A.; Ahmad, I.S.; Abbasi, A.; Cunningham, B.T.; Watkin, K.L. Cytotoxicity screening of Bangladeshi medicinal plant extracts on pancreatic cancer cells. *BMC Complement Altern Med.* **2010**, *17*, 10–52. <https://doi.org/10.1186/1472-6882-10-52>.
136. Hossain, A.I.; Faisal, M.; Rahman, S.; Jahan, R.; Rahmatullah, M. A preliminary evaluation of antihyperglycemic and analgesic activity of *Alternanthera sessilis* aerial parts. *BMC Complement Altern Med.* **2014**, *24*, 14–169. <https://doi.org/10.1186/1472-6882-14-169>.
137. Thomas, W.; Merish, S.; Tamizhamuthu, M. Review of *Alternanthera sessilis* with reference to traditional siddha medicine. *Int. J. Pharmacogn. Pharm. Res.* **2014**, *6*, 249–254.
138. Tene, V.; Malagón, O.; Finzi, P.V.; Vidari, G.; Armijos, C.; Zaragoza, T. An ethnobotanical survey of medicinal plants used in Loja and Zamora–Chinchipe, Ecuador. *J. Ethnopharmacol.* **2007**, *111*, 63–81. <https://doi.org/10.1016/j.jep.2006.10.032>.
139. Mosyakin, S.; Robertson, K.; Amaranthus, L. *Flora of North of Mexico (Magnoliophyta: Caryophyllidae, Part 1)*; Flora of North America Editorial Committee, Ed.; Oxford University Press: Oxford, UK, 2003; pp. 410–435.
140. Sarker, U.; Oba, S. Nutrients, minerals, pigments, phytochemicals, and radical scavenging activity in *Amaranthus blitum* leafy vegetables. *Sci. Rep.* **2020**, *10*, 3868. <https://doi.org/10.1038/s41598-020-59848-w>.
141. Abbasi, A.; Khan, M.; Shah, M.H.; Shah, M.; Pervez, A.; Ahmad, M. Ethnobotanical appraisal and cultural values of medicinally important wild edible vegetables of Lesser Himalayas–Pakistan. *J. Ethnobiol. Ethnomed.* **2013**, *9*, 66. <https://doi.org/10.1186/1746-4269-9-66>.
142. Nedelcheva, A. An ethnobotanical study on wild edible plants in Bulgaria. *Eurasian J. Biosci.* **2013**, *7*, 77–94. <https://doi.org/10.5053/ejobios.2013.7.0.10>.
143. Abdiniyazova, G.J.; Khojimatov, O.K. Medicinal plants of Karakalpakstan. *Int. J. Sci. Res.* **2017**, *6*, 2113–2116. <https://doi.org/10.21275/art20178256>.
144. Shakeri, A.; Hazeri, N.; Vlizadeh, J.; Ghasemi, A.; Tavallaei, F.Z. Phytochemical screening, antimicrobial and antioxidant activities of *Anabasis aphylla* extracts. *Kragujevac J. Sci.* **2012**, *34*, 71–78.
145. Lakhdari, W.; Dehliz, A.; Acheuk, F.; M'lik, R.; Hammi, H.; Doumandji-Mitiche, B.; Gheriani, S.; Berrekbia, M.; Guernit, K.; Chergui, S. Ethnobotanical study of some plants used in traditional medicine in the region of Oued Righ (Algerian Sahara). *J. Med. Plants Stud.* **2016**, *4*, 204–211.
146. Pei, Y.; Yang, Z.D.; Sheng, J. Chemical constituents of *Anabasis salsa*. *Chem. Nat. Compd.* **2014**, *50*, 957–958. <https://doi.org/10.1007/s10600-014-1132-4>.
147. DanHong, C.; Cheng, Y.; Yan, W. Seed germination characteristics of the desert subshrub *Atriplex cana* and its ecological significance. *Acta Petrol. Sin.* **2015**, *24*, 131–138.
148. Soliman, G.A.; Abd El Raheim, M. Antihyperglycemic, antihyperlipidemic and antioxidant effect of *Atriplex farinosa* and *Atriplex nummularia* in streptozotocin-induced diabetes in rats. *Bull. Env. Pharmacol. Life Sci.* **2015**, *4*, 10–18.
149. Jeong, H.; Kim, H.; Ju, E.; Kong, C.-S.; Seo, Y. Antioxidant effect of the halophyte *Atriplex gmelinii*. *KSBB J.* **2016**, *31*, 200–207. <https://doi.org/10.7841/ksbbj.2016.31.4.200>.
150. Lefèvre, I.; Marchal, G.; Meerts, P.; Corréal, E.; Lutts, S. Chloride salinity reduces cadmium accumulation by the Mediterranean halophyte species *Atriplex halimus* L. *Environ. Exp. Bot.* **2009**, *65*, 142–152. <https://doi.org/10.1016/j.envexpbot.2008.07.005>.
151. Manousaki, E.; Kalogerakis, N. Phytoextraction of Pb and Cd by the Mediterranean saltbush (*Atriplex halimus* L.): Metal uptake in relation to salinity. *Environ. Sci. Pollut. Res. Int.* **2009**, *16*, 844–854. <https://doi.org/10.1007/s11356-009-0224-3>.
152. Mardi, P.; Fallah Huseini, H.; Ahvazi, M.; Tavakoli-Far, B. Effects of *Atriplex hortensis* hydroalcoholic extract on phenylhydrazine induced hemolytic anemia in rat. *J. Med. Plants By-Prod.* **2022**, *11*, 37–41. <https://doi.org/10.22092/jmpb.2020.352222.1276>.
153. Ghasemi, P.; Momeni, M.; Bahmani, M. Ethnobotanical study of medicinal plants used by Kurd tribe in Dehloran and Abadan districts, Ilam province, Iran. *Afr. J. Tradit. Complement Altern. Med.* **2013**, *10*, 368–385. <https://doi.org/10.4314/ajtcam.v10i2.24>.
154. Matloub, A.A.; Hamed, M.A.; El Souda, S.S. Chemo-protective effect on hepato–renal toxicity and cytotoxic activity of lipoidal matter of *Atriplex lindleyi* Moq. *Int. J. Pharm. Pharm. Sci.* **2014**, *6*, 187–196.
155. Hanganu, D.; Olah, N.; Dubei, N.; Mărculescu, A. Phytochemical studies on *Chenopodium bonus-henricus* L. *Proc. Rom. Acad. Series B* **2008**, *3*, 287–292.
156. Mabberley, D.J. *The Plant Book*; Cambridge University Press: Cambridge, UK, 1993; p. 119.
157. Koziok, M.J. Chemical composition and nutritional evaluation of quinoa (*Chenopodium quinoa* Willd.). *Food Comp. Anal.* **1992**, *5*, 35–68.
158. Yuan, C.-G.; Huo, C.; Gui, B.; Liu, P.; Zhang, C. Green synthesis of silver nanoparticles using *Chenopodium aristatum* L. stem extract and their catalytic/antibacterial activities. *J. Cluster Sci.* **2016**, *28*, 1319–1333. <https://doi.org/10.1007/s10876-016-1147-z>.
159. Roriz, C.L.; Barros, L.; Carvalho, A.M.; Santos-Buelga, C.; Ferreira, I.C. *Pterospartum tridentatum*, *Gomphrena globosa* and *Cymbopogon citratus*. A phytochemical study focused on antioxidant compounds. *Food Res. Int.* **2014**, *62*, 684–693. <https://doi.org/10.1016/j.foodres.2014.04.036>. hdl:10198/10145.
160. Anaswara, M.R.; Suresha, B.S.; Balasubramanian, T.; Sushma, Y.C.; Jaseela, N.M. Medical importance of *Gomphrena globosa* – A systematic review. *Asian J. Pharm. Hea. Sci.* **2022**, *12*, 2718–2721. <https://doi.org/10.5530/ajphs.2022.12.18>.

161. Yaseen, T.; Slathia, D.; Farooq, I. Improving floral characteristics and yield of globe amaranth (*Gomphrena globosa* L.) through pinching and application of bio fertilizers and its impact on soil fertility. *J. Pharm. Innov.* **2022**, *11*, 425–430. <https://doi.org/10.13140/RG.2.2.19073.74087>.
162. Gannoun, S.; Mahfoudhi, A.; Flamini, G.; Helal, A.; Mighri, Z. Chemical composition and antimicrobial activities of Tunisian *Salsola vermiculata* L. *J. Chem. Pharm. Res.* **2016**, *8*, 1087–1092.
163. Ribera, A.; Bai, Y.; Wolters, A.M.A.; van Treuren, R.; Kik, C. A review on the genetic resources, domestication and breeding history of spinach (*Spinacia oleracea* L.). *Euphytica* **2020**, *216*, 48. <https://doi.org/10.1007/s10681-020-02585-y>.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.